

Fractional Statistics in terms of the r -Generalized Fibonacci Sequences

M. Rachidi^{a 1}, E.H. Saidi^a and J.Zerouaoui^{a,b 2}

(a) Faculté des Sciences, Lab/UFR-Physique des Hautes Energies,
Avenue Ibn Battouta B.P. 1014, Faculté des Sciences de Rabat, Morocco

(b) The Abdus Salam International Centre for Theoretical Physics,
34100, PO BOX586, 34100, Trieste, Italy.

Abstract

We develop the basis of the two dimensional generalized quantum statistical systems by using results on r -generalized Fibonacci sequences. According to the spin value s of the 2d-quasiparticles, we distinguish four classes of quantum statistical systems indexed by $s = 0, 1/2 : mod(1)$, $s = 1/M : mod(1)$, $s = n/M : mod(1)$ and $0 \leq s \leq 1 : mod(1)$. For quantum gases of quasiparticles with $s = 1/M : mod(1)$, $M \geq 2$, we show that the statistical weights densities ρ_M are given by the integer hierarchies of Fibonacci sequences. This is a remarkable result which envelopes naturally the Fermi and Bose statistics and may be thought of as an alternative way to the Haldane interpolating statistical method.

PACS number: 05.30.-d/11.10.-z/11.30.Ly.

2000 MSC : 40A05, 40A25

¹*Département de Mathématiques et Informatique, Faculté des Sciences de Rabat, B.P. 1014, Rabat, Morocco.*

²*Laboratoire de Physique Théorique et Appliquée LPTA, Faculté des Sciences de Kénitra, Morocco*

Contents

1	Introduction	3
2	Generalized QSM	5
3	Haldane Statistics	8
4	Gas of Quasiparticles of Spin $1/3$	9
4.1	Interpolating Statistics	10
4.2	Generalized Exclusion Statistics	11
5	Fibonacci Sequences Setting	12
6	More on FQSM	15
7	Conclusion	18

1 Introduction

Quantum elementary excitations carrying generalized values of the spin, to which we shall refer here after as quasi-particles, are involved in the description of many aspects of the critical behaviours of physical systems in two and two plus one dimensions [1,2]. In effective field models of fractional quantum Hall (FQH) liquids for instance [3], the elementary excitations carry fractional values of the spin and fractional electric charge Q . In this kind of systems, the spin angle θ is a topological quantity which is generated by a Bohm Aharonov effect and eventually by a Berry phase for systems with non zero space curvature geometries. For FQH hierarchical states of generic level n , typically described by a $U(1)^n$ CS gauge theory with a matrix coupling K of integer entries K_{ij} $1 \leq i, j \leq n$ and integers l_i , the spin angle $\frac{\theta}{2\pi}$, proportional to $s = \sum_{i,j=1}^n l_i (K^{-1})_{ij} l_j$, is not necessary half integer. The spin s takes rational values and in general real ones.

From the mathematical point of view, the parameter s may, roughly speaking, be thought of as the highest weight of the $SO(2)$ representations. As such there is no restriction on it and, a priori, can take any value. This parameter, together with the scaling dimension δ were at the basis of the spectacular developments made over the two last decades in 2d critical systems [4,5,6].

In $2d$ conformal field theory (CFT), objects carrying generalized values of the spin are elegantly described by world sheet (WS) vertex field operators type $V_\alpha(\phi) = \exp(i\alpha\phi)$; where ϕ describes one or a collection of $2d$ free scalars and α is a parameter related to the square root of the spin s of $V_\alpha(\phi)$ [7]. Quantum states with general values of the spin are also involved in the description of the continuous spectrum of the Liouville theory [8], in the integrable thermal deformation of minimal models[9] including the $\mathbb{Z}_N$ Potts ones [10] and those related with $U_q(sl_2)$ quantum group representations[11] especially the periodic ones [12].

In statistical mechanics of many bodies, quasi-particles carrying generalized spins have been subject to some interest in the past. In [13]; these quasi-particles have been interpreted as virtual particles interpolating between bosons and fermions. This is explicitly seen on the Haldane statistical weight $W(\alpha)$ giving the number of accessible states of a gas of N quasi-particles occupying a group of G states:

$$W(\alpha) = \frac{[G - (\alpha - 1)(N - 1)]!}{N![G - \alpha N - (1 - \alpha)]!} \quad (1)$$

In this equation, α is an interpolating parameter lying between zero and one describing respectively bosons and fermions. Later on we will give the exact relation between α and the spin s .

Starting from this statistical weight formula, Wu has studied in [14] the thermodynamic properties of gases of quasi-particles with exotic spins and derived the basic equations of the general form of the distribution and the grand partition function. These equations are exactly solved for bosons, fermions and for the so called semions. Some of the results of the Wu study were rederived in [15,16] by using a completely different manner based on an extension of the Pauli exclusion principle [17]. The key idea of this method may be summarized as follows. Considering a gas of quasi-particles of spin $1/M \bmod 1$, M integer greater than one ($M \geq 2$) and assuming that the quantum statistics of quasi-particles of spin $1/M$ are governed by a generalization of the Pauli exclusion principle according to which no more than $(M - 1)$ quasi-particles can live altogether on the same quantum state. This principle extends the usual Pauli principle for fermions, associated here to the special case $M = 2$. In [15,16], see also [18], it has been shown that one can solve exactly several statistical mechanics quantities for this kind of quantum gases. The obtained results are in complete agreement with the Wu analysis [14].

In this paper we reconsider the study of [15,16] by examining the basis of such kind of quantum statistical systems using results on combinatorial aspects of generalized r-Fibonacci sequences. Instead of the interpolation formula of Haldane eq(2), we propose a new statistical weight $\rho(N + G, M)$, given by the following combinatorial expression of M -generalized Fibonacci sequences:

$$\rho(n, M) = \sum_{g_0+2g_1+\dots+Mg_{M-1}=n-M} \frac{(g_0 + g_1 + \dots + g_{M-1})!}{g_0!g_1!\dots g_{M-1}!} p_0^{g_0} p_1^{g_1} \dots p_{M-1}^{g_{M-1}}, \quad (2)$$

where the p_i 's, $0 \leq p_i \leq 1$, are interpreted as occupation probabilities. To have an idea on how this relation applies in practice, it is instructive to consider the pure case where the p_i 's are set to one and so the above formula reduces to the well known M -generalized Fibonacci numbers namely:

$$\rho(n, M) = \sum_{g_0+2g_1+\dots+Mg_{M-1}=n-M} \frac{(g_0 + g_1 + \dots + g_{M-1})!}{g_0!g_1!\dots g_{M-1}!}. \quad (3)$$

Moreover setting $g_0 + g_1 + \dots + g_{M-1} = G$ or equivalently $g_1 + 2g_2 + \dots + (M - 1)g_{M-1} = N$, one obtains the generalized weight density of a gas of

quasi-particles of spin $1/M$, whose leading case $M = 2$, gives just the usual fermions statistical weight $W(1) = \rho(N + G, 2)_{|g_0+g_1=G} = \frac{G!}{N!(G-N)!}$.

This paper is organized as follows. In section 2, we give the set up of quantum statistics of quasi-particles carrying general values of the spin and fix terminologies. In section 3, we review some general features of the Haldane Statistics and compare it to our approach. In section 4, we make an explicit study of the quantum gas of quasi-particles of spin $s = 1/3$ either by using standard techniques or in terms of $M = 3$ -generalized Fibonacci numbers. In section 5 we review the M -generalized Fibonacci sequences and exhibit their connection with fractional statistics. In section 6 we make a discussion and give our conclusion.

2 Generalized QSM

Statistics is a quantum mechanical feature of identical particles which governs the quantum behaviour of a collection of N particles in the large limit of N . Usually one distinguishes two kinds of quantum statistical systems: bosons carrying integer values of the spin ($s = 0 \bmod 1$) and fermions carrying half odd integers spins ($s = 1/2 \bmod 1$). The thermodynamic statistical properties of bosons and fermions are described by two different quantum statistical distributions namely the Bose and Fermi densities. In this section we want to discuss the general set up of the quantum statistics of systems of quasi-particles carrying general values of the spin. Then we give some results regarding the quantum statistics of quasi-particles of spin $1/M$, $M \geq 2$.

To start recall that in two dimensions, there is no restriction on the values the spin s ; it can take any value; i.e: $0 \leq s \leq 1 \bmod 1$. Thus, according to the values of the spin s , the quasi-particles describing the quantum excitations in 2d systems may be classified in different kinds and so it is natural to conjecture the existence of various types of quantum statistical systems in two dimensions. These are given by the following classification:

- QSM; quantum statistical systems with quasi-particles carrying spins $s = 0, 1/2 \bmod 1$.
- FQSM; quantum statistical systems with quasi-particles carrying fractional values of s ; i.e: $s = 1/M \bmod 1$, where $M \geq 2$

- RQSM; quantum statistical systems with quasi-particles carrying spins s type: $s = n/M : \text{mod}(1)$, where $M \geq 2$ and $M \neq k.n$ with k an integer.
- GQSM; quantum statistical systems with quasi-particles carrying spins s , with $0 \leq s \leq 1 : \text{mod}(1)$.

This classification, according to the values of the spin s , obeys naturally to the following embedding:

$$QSM \subset FQSM \subset RQSM \subset GQSM. \quad (4)$$

It is not difficult to see that QSM, describing bosons ($s = 0 \text{ mod } (1)$) and fermions ($s = 1/2 \text{ mod } (1)$), is recovered as the boundary limit of FQSM by taking respectively $M = +\infty$ and $M = 2$. Similarly, FQSM may be viewed as a particular case of RQSM by setting $n = 1$. Finally, putting $s = n/M$ in GQSM by restricting to quasi-particles with rational values of the spin, one obtains RQSM.

From the above classification, we expect that several features of QSM may be extended iteratively to the other statistical systems. For example the well known fermionic number operator $N = [(-)^F]$ extends naturally as

$$\langle \psi | N | \psi \rangle = \langle \psi | \exp(i2\pi F) | \psi \rangle \quad (5)$$

$$= \exp(i2\pi s), \quad (6)$$

where $|\psi\rangle$ is the wave function of a given quasi-particle state of spin s . The eigenvalues of the operator F are effectively given by the spin of the $|\psi\rangle$ eigenfunctions; namely $s = 0 : \text{mod}(1)$ for bosons, $s = 1/2 : \text{mod}(1)$ for fermions, $s = 1/M : \text{mod}(1)$ for FQSM, $s = n/M : \text{mod}(1)$ for RQSM and $0 \leq s \leq 1 : \text{mod}(1)$ for GQSM.

An other example of features of QSM which extends naturally to the other quantum systems of eq(4) is the Pauli principle. This exclusion principle may be generalized to all other quantum statistical systems. This is easily seen by computing the trace of fermionic number operator $\exp(i2\pi F)$ on representation spaces involving all possible values of the spin; in particular for the supermultiplets built of states ($s = 0, s = 1/2$) for QSM and their generalizations ($s = j/M, : 0 \leq j \leq M - 1$) for FQSM.

$$\text{Tr} : \exp(i2\pi F) = 1 - 1 = 0, \quad \text{for } QSM, \quad (7)$$

$$Tr : expi(2\pi F) = \sum_{j=0}^{M-1} [expi(\frac{2\pi j}{M})] = 0, \quad \text{for } FQSM, \quad (8)$$

Such relation extends to RQSM and GSSM as well. For the supersymmetric multiplets ($s = jn/M, : 0 \leq j \leq M - 1$) of the RQSM systems; we have

$$Tr : expi(2\pi F) = \sum_{j=0}^{M-1} [expi(\frac{2\pi jn}{M})] = 0, \quad \text{for } RQSM, \quad (9)$$

while for the multiplets ($s = \beta, : 0 \leq \beta \leq 1$) of GQSM systems, we have the following formula:

$$Tr : expi(2\pi F) = \int_0^1 ds [expi(2\pi s)] = 0, \quad \text{for } GQSM. \quad (10)$$

Having classified the various kinds of quantum statistical systems, we turn now to describe briefly methods used in studying gases of quasi-particles. There are several ways of doing. The first way, due to Haldane, is based on the combinatorial formula eq(1) which turns out to be the simplest and the natural weight one may conjecture. This approach uses linear interpolating relations between the Bosons and the Fermions and concerns quasiparticles of spin s lying between $1/2$ and 1 . Haldane quasiparticles obey a generalized exclusion Pauli principle although a clear statement of this principle for the continuous values of s is still lacking. A second approach which was suggested in [19], follows the same line of reasoning as of Haldane; but instead of looking for interpolating statistical weight, one considers interpolating wave functions. In other words if we denote by ϕ and ψ respectively the Bose and Fermi wave functions of a system of N particles, then the wave function χ of gases of quasiparticles of spin s is, roughly speaking, given by:

$$\chi = \phi \cos(2\pi s) + \psi \sin(2\pi s); 0 \leq s \leq 1/2. \quad (11)$$

However, a detailed analysis of this eq shows that for the case of quasi-particles of spin $s = 1/M$ one discovers exactly the quasi-determinant of [15,16] describing the wave function of the FQSM gas of spin $s = 1/M \pmod{1}$; with $M \geq 2$.

A third way, which was developed concerns gases of quasiparticles of spin $s = 1/M \pmod{1}$, is based on the idea of imposing from the beginning the generalized Pauli exclusion principle according to which no more than $(M - 1)$ quasiparticles of spin $1/M$; $M \geq 2$ can live altogether on the same

quantum state. In this approach, one obtains explicit relations for FQSM extending the Bose and Fermi analogue; in particular the following quantum distributions

$$\langle n_k \rangle = \frac{\xi}{1 - \xi} - \frac{M\xi^M}{1 - \xi^M}, \quad (12)$$

where we have set $\xi = e^{-\beta(\epsilon_k - \mu)}$ and $M \geq 2$. This eq offers an alternative to the standard quantum q-uo-n one given by $\langle n(q) \rangle = \frac{\xi}{1 - q\xi}$; where $-1 \leq q \leq 1$ and where for $q = 1, -1, 0$ we get respectively the Bose, Fermi and Boltzman distributions. .

In the next section, we review the main lines of the Haldane statistics eq(1). Then, we develop further the alternative we considered in [15,16]. Using group theoretical methods and combinatorial analysis we show that the rigorous results we obtained are not fortuitous but are due to the existence of a link between FQSM systems and a well known subject in mathematical physics namely Fibonacci sequences hierarchies. To that purpose, we shall proceed as follows:

(a) first review the basis of the Haldane approach; this method is based on a linear interpolating formula between bosons and fermions. The simplicity of this way is due to the fact that the weight density is due to the remarkable properties of the special function $\beta(x, y)$. Non linear interpolations between bosons and fermions were also considered in [19]; they involve hypergeometric functions $\mathcal{F}(a, b; \alpha, \beta)$ and.

(b) Develop a new setting using rigorous analysis involving group theoretical techniques and combinatorial formulas and show that the obtained results are indeed linked to generalized Fibonacci sequences.

3 Haldane Statistics

We start this section by describing some general features of the Haldane statistics. Although exact solutions are still missing, this statistics exhibits many remarkable features which make it very special. Haldane statistics based on eq(1), corresponds in fact to a linear interpolating statistics between bosons and fermions. This linearity is a useful mathematical feature which make the interpolating formula natural and simple to handle. Indeed, setting

$$\begin{aligned} x &= x(\alpha) = N \\ y &= y(\alpha) = G - \alpha N + (\alpha - 1), \end{aligned} \quad (13)$$

in eq(1), one sees that it reduces to the inverse of the well known di-gamma special function $\beta(x, y)$; namely:

$$W(x, y) = \frac{(x+y)!}{x!y!}, \quad (14)$$

$$W(x, y) \cdot \beta(x, y) = 1. \quad (15)$$

Note that $y(\alpha)$ is a linear interpolation between the Bose point y_B and the Fermi one y_F respectively given by:

$$\begin{aligned} y_B &= y(1) = G - N \\ y_F &= y(0) = G - 1. \end{aligned} \quad (16)$$

The interpolation eq(13) as well as the statistics of the boundary points eq(16), shows that the Haldane interpolating parameter α and the spin s are related in the large N limit as

$$\alpha = \frac{N(1-2s)}{N-2} \sim (1-2s), 0 \leq s \leq \frac{1}{2}. \quad (17)$$

For $s = 0$, and $s = \frac{1}{2}$, the parameter α is equal to one or zero respectively. For the special case $\alpha = \frac{1}{2}$, corresponding to semions of spin $s = \frac{1}{4}$, the Haldane statistics can be solved exactly; one finds the following quantum distribution for semions is given by:

$$\langle n_j(\alpha = 1/2) \rangle = \frac{1}{[\frac{1}{4} + \exp 2\beta(\epsilon_j - \mu)]^{1/2}} \quad (18)$$

For general values of α , the situation is however complicated because one has to solve non linear eqs of the type:

$$\begin{aligned} \langle n_j \rangle &= \frac{1}{w(e^{\beta(\epsilon_j - \mu)}) + \alpha}, \\ \xi &= w(\xi)^\alpha [1 + w(\xi)]^{1-\alpha}. \end{aligned} \quad (19)$$

where $\xi = e^{\beta(\epsilon - \mu)}$, ϵ is the energy and μ is the chemical potential.

4 Gas of Quasiparticles of Spin 1/3

The gas of identical quasi-particles of spin 1/3 is the simplest system coming after the quantum Fermi gas. This is a quantum system of FQSM, which may be studied by using two routes:

(i) using the Haldane interpolating statistical weight $W(\alpha)$ eq(1) and fixing the parameter as $\alpha = \frac{N}{3(N-2)}$; i.e $s = 1/3$.

(ii) solving the constraint equations imposed by FQSM according to which no more than $(M - 1)$ quasi-particles of spin $s = 1/M$ can live altogether on the same quantum state.

4.1 Interpolating Statistics

Following Haldane and using eq(17), the statistical weight $W_{s=1/3}$ of a gas of N quasi-particles of $s = 1/3$ can be shown to be given by:

$$W_{s=1/3} = \frac{\Gamma\left[\frac{3(N-2)G-(N-1)(2N-6)}{3(N-2)}\right]\Gamma(N)}{\Gamma\left[\frac{3(N-2)G-2N^2-2N+6}{3(N-2)}\right]}. \quad (20)$$

This eq may be rewritten into an equivalent form as:

$$W_{s=1/3} = W(N, y), \quad (21)$$

where now $y = \frac{1}{3(N-2)}[3(N-2)G - N^2 + 6 - 2N]$. Having these relations at hand one can derive the thermodynamic statistical mechanical features of this gas of quasi-particles. For example, the mean occupation number $\langle n_k \rangle$ is obtained by solving eqs(19). For the case of a gas with pure occupation probabilities, the partition function $z(\xi)$ reads, at the statistical equilibrium, as:

$$z(\xi) = 1 + \xi + \xi^2. \quad (22)$$

Note that eq(22) lies between with the partition function $z_{Fermi}(\xi) = 1 + \xi$, and $z_{Bose}(\xi) = 1 + \sum_{n=1}^{\infty} \xi^n$ of individual fermions and bosons of quantum Fermi and Bose gases respectively. $z_{Fermi}(\xi)$ is given by just the sum of two terms as a consequence of the Pauli exclusion principle contrary to the bosonic gas where no exclusion rule exist. Therefore, the polynomial eq(22) of order 2 reflects clearly that quasiparticles of spin $s = 1/3$ obeys a generalization exclusion statistics principle according to which no more than two particles of spin $s = 1/3$ can live together on the same quantum state of the gas. This selection rule is the simplest exclusion statistics coming after the Pauli exclusion formula obeyed by fermions and constitutes the leading term of a more general hierarchical formula namely:

$$z(\xi) = 1 + \xi + \dots + \xi^{M-1}. \quad (23)$$

This relation extends eq(22) for the case of quasi-particles of spins $\frac{1}{M}$, $M \geq 2$. In what follows we will show that such relation may be also derived by using an other alternative using combinatorial analysis.

4.2 Generalized Exclusion Statistics

We start by examining the special exclusion statistics obeyed by quasi-particles with spin $s = 1/3$ by using group theoretical methods and combinatorial analysis. Then we work out the properties of the formula of the statistical weight at equilibrium one gets, in particular its relation to 3-generalized Fibonacci numbers. To do so, consider a system of N quasi-particles of spin $s = 1/M$ with $M = p + 1$, occupying G quantum states. On each state of the G ones, there can live altogether at most p quasi-particles as required by the generalized exclusion statistics for particles with spin $s = 1/(p + 1)$. To get the total number of accessible state of the $p = 2$ system, we proceed as follows: First divide the set of the allowed G quantum states into three kinds of sub-sets g_0, g_1 and g_2 consisting respectively of B_0, B_1 and B_2 boxes which can contain zero, one and two quasi-particles of spin $s = 1/3$ of the system with occupation probabilities p_0, p_1 and p_2 with $0 \leq p_i \leq 1$. Thus, we have the following equalities:

$$g_0 + g_1 + g_2 = G, \quad (24)$$

$$g_1 + 2g_2 = N. \quad (25)$$

Since the B_0, B_1 and B_2 boxes are indistinguishable, it follows that for a given configuration of the quantum system; the number of accessible states is no longer given by $G!p_0^{g_0}p_1^{g_1}p_2^{g_2}$ but rather by:

$$\frac{G!}{g_0!g_1!g_2!}p_0^{g_0}p_1^{g_1}p_2^{g_2}. \quad (26)$$

For configurations associated with pure occupation probabilities; i.e $p_0 = p_1 = p_2 = 1$, the above relation reduces to the well known $\frac{G!}{g_0!g_1!g_2!}$ factor. Taking into account all possible partitions of the number N , the total number $D_3(N, G)$ of accessible states of the quantum gas occupying G quantum states reads then as:

$$D_3(N, G) = \sum_{g_1, g_2 \geq 0, g_1 + 2g_2 = N} \frac{G!}{g_1!g_2!(G - g_1 - g_2)!}p_0^{g_0}p_1^{g_1}p_2^{g_2}, \quad (27)$$

where we have used eq(25). Furthermore solving g_1 in terms of N and g_2 by help of eq(26), one may rewrite this equation into following equivalent form.

$$D_3(N, G) = \sum_{g=0}^{[N/2]} \frac{G!}{g!(N-2g)!(G+g-N)!} p_0^{g_0} p_1^{g_1} p_2^{g_2}, \quad (28)$$

where we have set $g = g_2$ and where $[N/2]$ is the integer part of $N/2$. Note in passing that this eq has some remarkable features; one of which is that $D_3(G, N)$ contains the fermion statistical density $D_2(G, N)$ as the leading term of the sum over the possible partitions of the B_2 boxes. In other words; $D_3(N, G)$ may be written, for pure probabilities, as

$$D_3(N, G) = \frac{G!}{N!(G-N)!} + \sum_{g=1}^{[N/2]} \frac{G!}{g!(G-2g)!(G+g-N)!}, \quad (29)$$

where the first term of the right hand side (rhs) of is just the statistical weight $D_2(N, G)$ of N fermions occupying G states. The remaining $([N/2] - 1)$ extra terms of eq(30) describe then a deviation above the $D_2(G, N)$ density and so it may be interpreted as the effects of quasi-particles of spin $s = 1/3$. Put differently, we have

$$D_3(G, N) = D_2(G, N) + \Delta_{32}(G, N), \quad (30)$$

where $\Delta_{32} = D_3(G, N) - D_2(G, N)$, given by the second term of the rhs of eq(32), describes the manifestation of FQSM.

5 Fibonacci Sequences Setting

A careful observation of the previous combinatorial analysis shows that it has much to do with the combinatorial realization of the generalized Fibonacci sequences[20], see also[21,22]. To exhibit this connection explicitly, let us first give some elements on the combinatorial aspects of these sequences, in particular for the Fibonacci sequences of order $M=2$ and 3. To start let a_0, a_1 be some fixed real numbers with $a_1 \neq 0$ and consider the sequence $\{V_n\}_{n=0}^{+\infty}$ defined by $V_n = \rho(n+1, 2)$ with,

$$\rho(n, 2) = \sum_{g_0+2g_1=n-2} \frac{(g_0+g_1)!}{g_0!g_1!} a_0^{g_0} a_1^{g_1},, \text{ for every } n \geq 3, \quad (31)$$

and $\rho(2, 2) = 1$, $\rho(n, 2) = 0$ for $n = 0, : 1$. Using this eq it is not difficult to verify that $V_0 = 0$, $V_1 = 1$ and

$$V_{n+1} = a_0 V_n + a_1 V_{n-1}, \text{ for every } : n \geq 1, \quad (32)$$

implying in turns that $\{V_n\}_{n=0}^{+\infty}$ is nothing else a weighted Fibonacci sequence with initial conditions specified by the numbers V_0, V_1 and weights a_0, a_1 . Note that for $a_0 = a_1 = 1$, we recover the classical Fibonacci numbers,

$$\rho(n, 2) = \sum_{g_0+2g_1=n-2} \frac{(g_0+g_1)!}{g_0!g_1!}, , :: \text{ for every } :: n \geq 3. \quad (33)$$

Observe also that up on imposing the extra constraint $g_0 + g_1 = G$, the V_n is intimately related to the fermions number density $\mathcal{D}_2(G, N)$. Indeed putting the condition $g_0 + g_1 = G$ back into eq(36), one discovers that sum over g_1 and g_0 with $g_0 + 2g_1 = n - 2$ is completely fixed and so $n = N + G + 2$ as well as:

$$\mathcal{D}_2(G, N) = \rho(N + G + 2, 2)|_{g_0+g_1=G}, \quad (34)$$

showing that the fermion density $\mathcal{D}_2(G, N)$ is given by a restricted Fibonacci numbers sequence of rank two. An inspection of this result shows that it is not a fortuitous feature and may be used to describe not only fermions but also statistical weights of quantum gases of fractional spin $s = 1/M$. This observation is supported by several arguments, above all the fact that Fibonacci sequences admit integer hierarchies in perfect coherence with FQSM systems. To illustrate the idea, we consider in what follows the $M = 3$ case; i.e the case of a quantum gas of quasi-particles of spin $1/3$. We begin however by giving some generalities on 3-generalized Fibonacci sequences, then we derive the relation linking D_3 and $\rho(n, 3)$. Later on we give the general result. Let a_0, a_1, a_2 be some fixed real numbers with $a_2 \neq 0$. If $a_2 = 0$ but $a_1 \neq 0$, one falls in the previous case. A 3-generalized Fibonacci sequences $\{V_n\}_{n=0}^{+\infty}$ is roughly speaking defined by:

$$V_{n+1} = a_0 V_n + a_1 V_{n-1} + a_2 V_{n-2}, \text{ for every } : n \geq 2, \quad (35)$$

where V_0, V_1, V_2 and a_0, a_1, a_2 are respectively the initial conditions and the weights of $\{V_n\}_{n=0}^{+\infty}$. Eq(38) extends the usual Fibonacci sequences $\{V_n\}_{n=0}^{+\infty}$ obeying eq(44), and which may also baptized as the rank 2 Fibonacci sequences; the leading element of the Fibonacci hierarchy. Recall by the

way that r-generalized Fibonacci sequences, which have been studied by using various method and techniques, go beyond eqs(44) and (47), and play a fundamental role in many fields of applied sciences and engineering []. In practice there are two ways to handle these sequences: the combinatorial realization which we consider hereafter and the Binet representation. For the third order of the hierarchy we are discussing, the combinatorial realization is generated by:

$$\rho(n, 3) = \sum_{g_0+2g_1+3g_2=n-3} \frac{(g_0 + g_1 + g_2)!}{g_0!g_1!g_2!} a_0^{g_0} a_1^{g_1} a_2^{g_2}, \text{ for every } n \geq 3, \quad (36)$$

where we have set $\rho(3, 3) = 1$ and $\rho(j, 3) = 0$ for $0 \leq j \leq 2$. To check that the above eq is indeed a 3-generalized Fibonacci sequence, it is enough to take $V_n = \rho(n + 1, 3)$. Straightforward calculations show that we have $V_0 = V_1 = 0$ and $V_2 = 1$ as well as:

$$V_{n+1} = a_0 V_n + a_1 V_{n-1} + a_2 V_{n-2}, \text{ for every } n \geq 2.$$

Note in passing that for $a_0 = a_1 = a_2 = 1$, the sequence $\{V_n\}_{n=0}^{+\infty}$ gives the well known 3-generalized Fibonacci numbers, very familiar in the mathematical literature[M],[K]. In this case, eq (45) becomes:

$$\rho(n, 3) = \sum_{g_0+2g_1+3g_2=n-3} \frac{(g_0 + g_1 + g_2)!}{g_0!g_1!g_2!}, \text{ for every } n \geq 3, \quad (37)$$

with $\rho(3, 3) = 1$ and $\rho(j, 3) = 0$ for $0 \leq j \leq 2$. The beauty of this relation is that it is linked to the density of the quantum gas of quasi-particles of spin $s = 1/3$ we studied earlier. Indeed, if instead of the two constraint eqs (17-18), we consider rather the following relaxed condition

$$g_0 + 2g_1 + 3g_2 = G + N, \quad (38)$$

obtained by adding eq(17) and eq(18), then the density $D_3(G, N)$, given by eqs(28-29), is translated exactly to eq(46). Put differently $D_3(G, N)$ can be derived from the 3-generalized Fibonacci numbers by imposing either $g_1 + 2g_2 = N$ or $g_0 + g_1 + g_2 = G$. In the formulation based on the Fibonacci sequences, the number of accessible states eq(19) is now replaced by:

$$\frac{(g_0 + g_1 + g_2)!}{g_0!g_1!g_2!} p_0^{g_0} p_1^{g_1} p_2^{g_2}. \quad (39)$$

Taking into account of all possible configurations of the number $N + G$ partitions in eq(41), the new total number $\rho_3(N, G) = \rho(G + N - 3, 3)$ of accessible states is:

$$\rho_3(N, G) = \sum_{g_0+2g_1+3g_2=G+N} \frac{(g_0 + g_1 + g_2)!}{g_0!g_1!g_2!} p_0^{g_0} p_1^{g_1} p_2^{g_2}, \quad (40)$$

which could be compared with eq(39). Introducing the Dirac delta function $\delta(x)$, we can write,

$$D_3(G, N) = \delta(g_0 + g_1 + g_2 - G) \rho(G + N - 3, 3). \quad (41)$$

Substituting the delta function and solving g_1 in terms of N and g_2 , we can rewrite eq(20) into the following equivalent form:

$$D_3(N, G) = \sum_{g=0}^{[N/2]} \frac{G!}{g!(N-2g)!(G+g-N)!}, \quad (42)$$

where we have set $g = g_2$ and where $[N/2]$ is the integer part of $N/2$. Having this relation at hand, we can write down all thermodynamics properties of the gas of quasi-particles at equilibrium.

6 More on FQSM

In this section we want to extend some of the previous results to quantum systems of N quasi-particles of spin $1/M$; $M \geq 2$ using the combinatorial M-generalized Fibonacci sequences. We start by generalizing the useful properties on the combinatorial and the Binet expressions for the M-generalized Fibonacci sequences.

Let $a_0, a_1, \dots, a_{M-1}$ be $(M-1)$ arbitrary real numbers with $a_{M-1} \neq 0$ and $M \geq 2$. An M-generalized Fibonacci sequence $\{V_n\}_{n=0}^{+\infty}$ is defined as follows:

$$V_{n+1} = a_0 V_n + a_1 V_{n-1} + \dots + a_{M-1} V_{n-M+1}, \text{ for } n \geq M-1 \quad (43)$$

where $V_0, V_1, \dots, V_{M-1}$ and $a_0, a_1, \dots, a_{M-1}$ are the initial conditions the sequence. In case where $a_i = 1$ for all i 's and $V_0 = V_1 = \dots = V_{M-2} = 0$ and $V_{M-1} = 1$, we recover the M-generalized Fibonacci numbers while for $a_{M-1} = 0$ and $a_{M-2} \neq 0$, one gets $(M-1)$ -generalized Fibonacci sequences

to which one can apply the same analysis. The combinatorial expression of $\{V_n\}_{n=0}^{+\infty}$, for $n \geq M$, is realized as:

$$\rho(n, M) = \sum_{g_0+2g_1+\dots+Mg_{M-1}=n-M} \frac{(g_0 + g_1 + \dots + g_{M-1})!}{g_0!g_1!\dots g_{M-1}!} a_0^{g_0} a_1^{g_1} \dots a_{M-1}^{g_{M-1}}, \quad (44)$$

where we set $\rho(M, M) = 1$ and $\rho(n, M) = 0$ for $0 \leq n \leq M - 1$. Starting from the following identity

$$\frac{(k_0 + \dots + k_{M-1})!}{k_0!k_1!\dots k_{M-1}!} = \sum_{j=0}^{M-1} \frac{(k_0 + \dots + k_{M-1} - 1)!}{k_0!\dots k_{j-1}!(k_j - 1)!k_{j+1}\dots k_{M-1}!}$$

we derive easily that,

$$\rho(n+1, M) = a_0\rho(n, M) + a_1\rho(n-1, M) + \dots + a_{M-1}\rho(n-M+1, M),$$

Thus the sequence $\{W_n\}_{n=0}^{+\infty}$ defined by $W_n = \rho(n+1, M)$, is a sequence whose initial conditions are such that $W_0 = W_1 = \dots = W_{M-2} = 0$ and $W_{M-1} = 1$. More generally, the combinatorial expression of the sequence (46) can be given, for every $n \geq M$, in terms of $\rho(n, M)$ (47) as follows.

$$V_n = A_0\rho(n, M) + A_1\rho(n-1, M) + \dots + A_{M-1}\rho(n-M+1, M), \quad (45)$$

where

$$A_m = a_{M-1}V_m + a_{M-2}V_{m+1} + \dots + a_mV_{M-1}; \text{ for } m = 0, 1, \dots, M-1,$$

and where $\rho(M, M) = 1$ and $\rho(j, M) = 0$ for $j < M$.

In the Binet representation of M -generalized Fibonacci sequence, the V'_n s are expressed as polynomials in the eigenvalues of:

$$P(X) = X^M - a_0X^{M-1} - \dots - a_{M-2}X - a_{M-1}, \quad (46)$$

The point is that to every sequence $\{V_n\}_{n=0}^{+\infty}$ of rank M , we associate the polynomial $P(X)$ above

whose roots $\lambda_0, \lambda_1, \dots, \lambda_s$ have respectively the multiplicities $m_0, m_1, \dots, m_s$. The Binet realisation of the general term V_n of the sequence reads in terms of the λ'_i s as:

$$V_n = \sum_{l=1}^s \sum_{j=0}^{m_s-1} \beta_{l,j} n^j \lambda_l^n, \quad (47)$$

where the $\beta'_{l,j}$ s are obtained from initial conditions by solving the following system of M linear equations

$$\sum_{l=1}^s \sum_{j=0}^{m_s-1} \beta_{l,j} n^j \lambda_l^n = V_n, \text{ for } n = 0, 1, \dots, M-1. \quad (48)$$

Such expressions are known in the mathematical literature as the Binet formula of $\{V_n\}_{n \geq 0}$. For the leading $n=2$ case where the λ_j solutions of the characteristic polynomial $P(x) = x^2 - a_0x - a_1$ have are simple to handle, we have

$$V_n = \sum_{j=1}^2 B_j \lambda_j^n, \text{ for } \lambda_1 \neq \lambda_2, \quad (49)$$

$$V_n = (B_1 + B_2 n) \lambda_1^n, \text{ for } \lambda_1 = \lambda_2, \quad (50)$$

where the B_j 's are related to α_0 and α_1 as

$$\sum_{j=1}^2 B_j \lambda_j^n = \alpha_n; \quad n = 0, 1 \text{ (for } \lambda_1 \neq \lambda_2) \quad (51)$$

$$(B_1 + B_2 n) \lambda_1^n = \alpha_n, \quad n = 0, 1 \text{ (for } \lambda_1 = \lambda_2). \quad (52)$$

Such eqs may written in a more explicit way by specifying the explicit form of the λ_j 's in terms of the initial parameters.

Having given some tools on Fibonacci sequences hierarchies, let us turn now to expose briefly the link with FQSM systems. To that purpose, we reconsider the gas described in section 4 but now the quasi-particles have spin $s = 1/M$ with $M = P + 1$, $P \geq 1$. In this case eqs(27-28) extend as

$$g_0 + g_1 + g_2 + \dots + g_P = G \quad (53)$$

$$g_1 + 2g_2 + 3g_3 + \dots + P g_P = N \quad (54)$$

The density of accessible states of the gas is,

$$\mathcal{D}_{P+1}(N, G) = \sum_{\substack{g_1, g_2, \dots, g_P \geq 0 \\ \sum_{j=1}^P j \cdot g_j = N}} G! \left[\left(G - \sum_{k=1}^P G_P \right)! \prod_{l=1}^P (g_l!) \right]^{-1} \quad (55)$$

Eq(37) exhibits several features, some of them extends properties of the $\mathcal{D}_3(N, G)$ statistical weight discussed earlier. This concerns for instance the

interpretation of $\mathcal{D}_{P+1}(N, G)$ as a deviation above the $\mathcal{D}_P(N, G)$ weight of quasi-particles of spin $1/P$. Indeed rewriting eq(54) by singling out the sum over g_P ; i.e,

$$\mathcal{D}_{P+1}(N, G) = \sum_{g_P=0}^{[N/P]} \frac{1}{g_P!} \sum_{\substack{g_1, g_2, \dots, g_{P-1} \geq 0 \\ \sum_{j=1}^{P-1} j \cdot g_j = N - P g_P}} \frac{G!}{g_1! g_2! \dots, g_{P-1}! (G - \sum_{j=1}^P g_j)!} \quad (56)$$

one sees that the leading term in the sum over g_P ; ie for $g_P = 0$, is just the weight density $\mathcal{D}_P(N, G)$ of a gas of N quasi-particles of spin $1/P \pmod{P}$. Put differently, eq(55) may usually decomposed as

$$\mathcal{D}_{P+1}(N, G) = \mathcal{D}_P(N, G) + \Delta_{P+1, P}; \quad (57)$$

where $\Delta_{P+1, P}$ measures the derivation from $\mathcal{D}_{P+1}(N, G)$ to $\mathcal{D}_P(N, G)$ densities and reads as

$$\Delta_{P+1, P} = \sum_{g_P=1}^{[N/P]} \frac{1}{g_P!} \sum_{\substack{g_1, g_2, \dots, g_{P-1} \geq 0 \\ \sum_{j=1}^{P-1} j \cdot g_j = N - P g_P}} \frac{G!}{g_1! g_2! \dots, g_{P-1}! (G - \sum_{j=1}^P g_j)!} \quad (58)$$

This eq is just a generalization of eqs(32-33); it is valid for any value of P greater than three and obeys quite remarkable properties inherited from the generalized Fibonacci combinatorial method we have been using. Eq(57) may be also thought as a manifestation of the exclusion statistics in FQSM stating that no more than P quasi-particles of spin $\frac{1}{P+1}$ can live altogether on the same quantum state. It satisfies equally the following splitting

$$\mathcal{D}_{p+1}(N, G) = [\mathcal{D}_2(N, G) + \sum_{j=3}^{p-1} \mathcal{D}_j(N, G)] + \sum_{j=2}^p \Delta_{j+1, j}; \quad (59)$$

in agreement with the embedding (4). Here the $\Delta_{j+1, j}$'s, given by similar relations to (57), are deviations from D_{j+1} to D_j densities.

7 Conclusion

In this paper we have developed a new way to approach generalized quantum statistical mechanics of gases of quasi-particles. Our method, which

is based on group theoretical analysis and combinatorial representation of M-generalized Fibonacci sequences, can be viewed as an alternative to the Haldane approach. To do so, we have first reviewed the setting of generalized quantum statistics which we have classified into four classes according to the values of the spin s of the quasi-particles: (1) the usual QSM for $s = 0, 1/2 \bmod(1)$, (2) FQSM for $s = 1/M \bmod(1)$, (3) RQSM for $s = n/M \bmod(1)$ and (4) GQSM for $0 \leq s \leq 1 : \bmod(1)$. This classification obeys the inclusions: $QSM \subset FQSM \subset RQSM \subset GQSM$.

Then we have reviewed the interpolating analysis of Haldane based on eq(1) and showed that it is equivalent to the method we developed in [14,15] using a generalization of the Pauli exclusion rule as a general principle.

Next motivated by a set of observations; in particular the embedding eq(4), we have developed an other way to deal with FQSM gases. Our method is based on the following weight density:

$$\rho(n, M) = \sum_{g_0+2g_1+\dots+Mg_{M-1}=n-M} \frac{(g_0 + g_1 + \dots + g_{M-1})!}{g_0!g_1!\dots g_{M-1}!} p_0^{g_0} p_1^{g_1} \dots p_{M-1}^{g_{M-1}}, \quad (60)$$

where the p_i 's are dilution probabilities. Up on imposing eqs(24-25) or equivalently eq(38), we rederive the known results. We believe that our way of doing offer a powerful manner to deal with such quantum systems. Our motivations for this claim are:

(1) Eq(60) is obtained by using a rigorous analysis based on a group theoretical method and a combinatorial analysis.

(2) Eq(60) is in agreement with the embedding $QSM \subset FQSM$. It recovers standard results of QSM and describes naturally any kind of quantum gas of quasi particles of spin $s = 1/M$, $M \geq 2$.

(3) Eq(60) is linked to a well known subject in mathematical physics; namely the combinatorial aspect of the M-generalized Fibonacci numbers. In this setting QSM is described by the standard Fibonacci number sequences while FQSM is described by its integer hierarchies. QSM appears then as the leading term of the generalized Fibonacci hierarchy. In this context we expect that RQSM and GQSM would be also described by means of similar analysis.

(4) Finally eq(60) incorporates in a natural way the dilution property of quantum gases carried by the p_i parameters. Therefore diluted quantum gases of FQSM are described by integer hierarchies of Fibonacci sequences.

Acknowledgment

One of the authors; J.Z. would like to thank the AS-ICTP and the High Energy Section for hospitality. He would also to acknowledge the Student program at ICTP for invitation.

This research work has been supported by the program PARS number: 372-98 CNR.

References

- [1] Daniel Friedan, Zong-an Qiu, Stephen Shenker, Phys.Rev.Lett.52:1575-1578,1984 Phys.Lett.B151:37-43,1985
C.Itzykson, J.M Drouffe, Statistical Field Theory (1989) Cambridge, UK: Univ. Pr (1989).
F.Wilczek, Fractional Statistics and Anyon supraconductivity, WS Singapore(1990).
- [2] R.B Laughlin, Quantum Hall Effect, R.E Prange and S.M Girvin (Eds).
- [3] X.G. Wen, Y.S. Wu and Y. Hatsugai, Nucl.Phys.B422:476-494,1994
Xiao-Gang Wen, Topological orders and Edge Excitations in FQH States,PRINT-95-148 (MIT).
A. Jellal Int.J.Theor.Phys.37:2751-2755,1998
I. Benkaddour, A. El Rhalami, E.H. Saidi; Non Trivial extension of the (1+2)-Poincare algebra and conformal Invariance on the Boundary of AdS_3 ; hep-th/0007142, To appear in Euro Jour C (2001).
- [4] A.A. Belavin, A.M Polyakov and A.B Zamolodchikov, Nucl.Phys.B241:333-380,1984.
- [5] A.B.Zamolodchikov, Rev.Math.Phys.1:197-234,1990. Adv.Stud.Pure Math.19:641-674,1989
- [6] J.Cardy, Conformal Invariance and Universality in Finite size Scaling, J.PHYS. A17 (1984) L385-L387.
- [7] Oleg Andreev, Boris Feigin Nucl.Phys.B433:685-711,1995,
P. Di Vecchia, M. Frau, A. Lerda and S.Sciuto Nucl.Phys.B298:527,1988.
- [8] Andre LeClair and Dennis Nemeschansky, Mod.Phys.Lett.A11:139-146,1996.
Denis Bernard and Andre Leclair, Nucl.Phys.B399:709-748,1993.
- [9] A.B Zamolodchikov, Nucl.Phys.B366:122-134,1991
G.Mussardo Phys Report 218 (1992)215.
- [10] A.Kadiri, E. H.Saidi, M. B.Sedra and J.Zerouaoui, ICTP preprint IC/94/216.
- [11] A.Leclair and C.Vafa Nucl Phys B401 (1993)413

- [12]E.H.Saidi, M.B.Sedra and J.Zerouaoui, Class. Quant.Grav. 12 (1995) 1576.
E.H.Saidi, M.B.Sedra and J.Zerouaoui, Class. Quant. Grav. 12 (1995) 2705.
- [13]F. D. H. Haldane, Phys. Rev. Lett. 67 (1991) 37.
- [14]Y. S. Wu, Phys. Rev. Lett. 73, N7 (1994).
- [15]E. H. Saidi and J. Zerouaoui, Revue Marocaine des Sciences Physique N2 (2000).
- [16]M. Rachidi, E. H. Saidi and J. Zerouaoui, Phys. Lett. B409 (1997) 394.
- [17]E. H. Saidi, ICTP preprint IC/95/299.
E. H. Saidi, Proceeding de la journee de Physique Statistique; Avril (1995), FST Settat, Morocco.
- [18]J. Zerouaoui, Doctorat d'Etat, Rabat Morocco (2000).
- [19] N.Fahssi, Lab/UFR-PHE (2000)Unpublished.
- [20]T. P. Dence, Ratios of generalized Fibonacci sequences. Fibonacci Quartely 25 (1987), 137-143. F. Dubeau, On 2-generalized Fibonacci sequences. Fibonacci Quartely 27 (1989), 221-229.
E. P. Miles. Generalized Fibonacci Numbers and associated Matrices, Amer.Math.Montly 67 (1960), 745-747.
- [21]M.Mouline and M.Rachidi, Suites de Fibonacci generalises et Chaines de Markov,
Revista de la Real Academia de Ciencias Exatos, Fisicas y Naturales de Madrid 89.1-2 (1995) 61-77.
- [22]F.Dubeau, W.Motta, M.Rachidi and O.Sachi, On Weighted r-GFS, The Fibonacci Quartely 35.2 (1997)102-110.
C.E Chidume, M.Rachidi and E.H Zerouali, Solving the general truncated Moment Problem by r-GFS method.
Journal of Mathematical Analysis and Applications **256(2001)625-635**.